



Using ComBase Predictor and Pathogen Modeling Program as support tools in outbreak investigation: an example from Denmark

Møller, Cleide; Hansen, Tina Beck; Andersen, Jens Kirk

Publication date:
2009

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Møller, C., Hansen, T. B., & Andersen, J. K. (2009). *Using ComBase Predictor and Pathogen Modeling Program as support tools in outbreak investigation: an example from Denmark*. Poster session presented at 6th International Conference on Predictive Modelling in Foods, Washington, DC, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Using ComBase Predictor and Pathogen Modeling Program as support tools in outbreak investigation: an example from Denmark



Tina Beck Hansen, Cleide O.A. Møller and Jens Kirk Andersen

National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark



QUESTION:

Could the *Yersinia enterocolitica*, found in children suffering from VTEC O26:H11 infection, also have come from the outbreak sausage?


METHOD:

- Step 1.** Collecting information on intrinsic factors of sausage
- Step 2.** Comparing growth & reduction rates using CBP & PMP
- Step 3.** Predicting growth & reduction during sausage production

ANSWER:

YES! *Yersinia enterocolitica* can grow in the sausage – according to ComBase Predictor and Pathogen Modeling Program.

Step 3. Results



- Growth of both *Y. enterocolitica* as well as VTEC was possible during the fermentation and the first part of drying
- Further growth of *Y. enterocolitica* was possible in the end of drying
- Slow reduction of VTEC but growth of *Y. enterocolitica* was possible during storage
- Fermentation error could have led to an additional 1 log increase of both *Y. enterocolitica* as well as VTEC at the point of release of the outbreak sausage

Reduction in PMP 7.0

Time to 1 log decrease =
time to 5 log reduction / 5

Growth in CBP & PMP 7.0

Time to 1 log increase =
doubling time · ln(10) / ln(2)

pH

Batter pH =

pH of raw meat → pH = 5.5

Sausage pH =

Dextrose added to boost the fermentation → pH = 4.8

Nutrients

Per 100 g sausage:

Fat 9 g (F)

Protein 22 g (P)

Carbohydrates 2 g (C)

List of ingredients

For 100 g (Y) sausage:

127 g meat (M), 4 g salt (N), 2 g spices (S), 0.5 g dextrose (D), no nitrite.

Step 2. Results

Growth / reduction rates during sausage processing

Step in processing	pH	WPS (w/v %)	Temperature (°C)	Time to 1 log change (without lag)	
				<i>Yersinia enterocolitica</i> From CBP	VTEC From PMP
Start fermentation	5.5	3.4	24	7.1 h	6.3 h
End fermentation	4.8	3.4	24	11 h	10 h
Start drying	4.8	3.4	16	25 h	27 h
Mid drying	4.8	5.0	16	45 h	40 h
End drying	4.8	6.0	16	3 d	8 d
Chilled storage	4.8	6.0	5	15 d	26 d

Step 1. Results

Water and salt contents in batter and ready-to-eat sausage

Calculation of	Formula	Result
Weight reduction (%)	$WR = 100 \cdot ((M+N+S+D) - 100) / (M+N+S+D)$	25 %
Salt content in batter (w/w %)	$Nb = 100 \cdot N / (M+N+S+D)$	3.0 w/w %
Water content in batter (%)	$Wb = 100 \cdot Ws / (100 - WR)$	84 %
Water phase salt in batter (w/v %)	$WPSb = 100 \cdot Nb / (Wb + Nb)$	3.4 w/v %
Salt content in ready-to-eat sausage (w/w %)	$Ns = 100 \cdot N / Y$	4.0 w/w %
Water content in ready-to-eat sausage (%)	$Ws = 100 - (F+P+C+Ns)$	63 %
Water phase salt in ready-to-eat sausage (w/v %)	$WPSs = 100 \cdot Ns / (Ws + Ns)$	6.0 w/v %